



BORMAX FOLIAREL 150 EC

FertiGlobal®
www.fertiglobal.it




Società Chimica Larderello spa
20124 Milano (Italy) - Via Fara 28
Tel +39 02 6771681 - Fax +39 0267716820/30

COMPANY WITH INTEGRATED
MANAGEMENT SYSTEM
CERTIFIED BY DNV
ISO 9001/2000
ISO 14001

10,9 B

EC FERTILISER
Boron ethanolamine

Composition	Water soluble boron (B)	10,9% (150 g/L)
Formulation	Solution (specific weight: 1,375 kg/L at 20°C)	
Packaging	12 x 1 L (12x1,375 kg) - 10 L (13,75 kg)	

Caution

In greenhouse do not apply concentrations above 100 mL/100 L. In case of mixture with other products try the compatibility beforehand on small surfaces.

Do not treat during flowering. PVA dispersible bags must be fully dissolved before adding FOLIAREL® Flüssig.

Carefully rinse the sprayer after use.

Crops	Rates		Application time & mode
	L/ha	L/100 L	
Sugar beet	2 - 4	0,8 - 1	At 8 and 16 leaf stage
Oilseed rape	2 - 4	0,8 - 1	1 application at leaf development and at stem elongation
Sunflower	2 - 4	0,8 - 1	1-2 application at 4-8 leaf stage
Legume grain crops	1	0,3 - 0,4	With plants 10-15 cm high and before flowering
Lucerne and Legume forage crops	1 - 3	0,5 - 0,8	After each cutting
Horticultural crops	1	0,3 - 0,4	After 4-6 leaves
Grapevines	1 - 3	0,2 - 0,3	Application at sprouting, visible bunch, fruit setting, berry development
Fruit trees	1 - 2	0,1 - 0,15	At fruit setting and post-harvest
Apple and Olive trees	2 - 3	0,1 - 0,2	Before flowering (2 applications) and at fruit settings
Flowers and ornamental plants	10 mL/100 m ²		Early growth
Fertigation	3 - 6		